


## A case of COVID-19 vaccination-associated *forme fruste* purpura fulminans

DOI: 10.1111/bjd.20744



DEAR EDITOR, A 77-year-old female patient developed multiple painful lesions 2 weeks after her first COVID-19 vaccination with mRNA-1273 (Moderna, Cambridge, MA, USA). (a) The figure shows the clinical presentation 4 weeks after the first COVID-19 vaccination (arrows indicate the distribution of lesions on the back). (b) Histopathology showed full-thickness necrosis of the dermis with thrombotic occlusion of virtually all capillaries and arterioles without concomitant inflammation suspicious of a septic embolic process. Coagulation occurred without pathology;<sup>1</sup> no relevant medication history was reported. (c) Two weeks later (6 weeks after vaccination), lesions were sharply demarcated (without treatment). Time kinetics, distribution of lesions and exclusion of other known causes<sup>2</sup> suggest that the purpura fulminans-like lesions were vaccination-associated. However, clinicians should not be diverted from careful investigation of other causes if they encounter similar events.

J. Griss ,<sup>1</sup> S. Eichinger ,<sup>2</sup> S. Winkler ,<sup>3</sup>  
W. Weninger <sup>1</sup> and P. Petzelbauer <sup>1</sup>

<sup>1</sup>Department of Dermatology, <sup>2</sup>Division of Hematology and Hemostaseology and <sup>3</sup>Division of Infectious Diseases and Tropical Medicine, Department of Medicine 1, Medical University of Vienna, Währinger Gürtel 18-20, Vienna, 1090, Austria

Email: johannes.griss@meduniwien.ac.at

### References

- 1 Greinacher A, Thiele T, Warkentin TE et al. Thrombotic thrombocytopenia after ChAdOx1 nCov-19 vaccination. *N Engl J Med* 2021; **384**:2092–101.
- 2 Chalmers E, Cooper P, Forman K et al. Purpura fulminans: recognition, diagnosis and management. *Arch Dis Child* 2011; **96**: 1066–71.

Funding sources: none.

Conflicts of interest: the authors declare they have no conflicts of interest.